

The Equivalence Principle and gravitational and inertial mass relation of classical charged particle

Mario Goto

*Departamento de Física/CCE
Universidade Estadual de Londrina
86051-990 Londrina, PR - Brazil
(mgoto@uel.br)*

Abstract

We show that the locally constant force necessary to get a stable hyperbolic motion regime for classical charged particles, actually, is a subtle combination of an applied external force and the radiation reaction force. It suggests, as the Equivalence Principle is valid, that the gravitational mass of charged particle should be slight greater than its inertial mass. However, an interesting new feature emerges from the unexpected behavior of the gravitational and inertial mass relation at very strong gravitational field, just reachable at the early stage of the universe or, perhaps, near massive black holes, with a divergence that introduces a critical field value. It signs that the Equivalence Principle should not be valid near to this critical field, which certainly coincides where the quantum effects turns to be relevant. We are using the Lorentz-Dirac equation to introduce the radiation reaction force and, as the Equivalence Principle is one of the foundations of the Einstein General Relativity, no reference to Einstein equations is made at any moment to develop this work, in order to avoid any vicious causal recurrence.

PACS: 03.50.D

I. INTRODUCTION

There is a old classical problem that survived unsolved until now with a quite general belief that it has no enough importance to be worthy to spend time trying to understand it, sometimes claimed as unsolvable without a quantal treatment ^{[17], [18]}. The problem in focus is the effect of the radiation reaction force on the charged particle dynamic, as given by the Lorentz-Dirac equation ^{[1]- [12]}, which should affect the validity of the Equivalence Principle at some circumstances ^{[9], [10], [21]}.

Perhaps it is better to believe that any problem that survive such a long time has to be important and it is to be worthwhile to spend any time as necessary to figure out where we are wrong. In accordance to this belief, there is now a renewed interest on this subject, with works pointing to something new ^{[20]- [22]}. By the way, we have to notice that some different features are emerging from radiation reaction force problem ^[19].

There are a lot of works about this subject accumulated since the first attempt was made by Dirac^[1]. However, the goal of this work is not to discuss about these papers, but, instead, to add a possibility to analyze this problem in a different perspective, with emphasis in the Equivalence Principle, which validity is used as a good starting point. From this point of view, one can show that what seems to be uncomfortable, as the presence of radiation for charged particle performing hyperbolic motion and its absence for one supported at rest in an uniform gravitational field^{[14], [15]}, both equivalent as the Equivalence Principle is valid, should lead to a new physical feature performed by charged particles. As the subject of this work is about the conditions of validity or not of the Equivalence Principle, which is one of the foundations of the Einstein General Relativity, it is just to notice that results from General Relativity are not used at any moment, in order to avoid any possibility to fall in a vicious causal recurrence.

What we have to do is to figure out the condition we have to provide such that a charged particle can reach a stable hyperbolic motion. It is necessary to furnish a subtle balance between an applied external force and the radiation reaction force to get an hyperbolic motion regime. An important consequence is that, taking account the Equivalence Principle, it implies a gravitational mass that is slight greater than the inertial mass. Perhaps because the main experimental justification that led Einstein to formulate the Equivalence Principle, which is one of the foundations of his General Theory of Relativity^[8], is the numerical equality between inertial and gravitational mass, nowadays they are taken quite as synonymous, so we have to be aware to avoid misleading conclusions. It is very important to emphasize the central role played by the radiation reaction force, so a slight upper deviation of gravitational mass should be necessary to validate the Equivalence Principle.

However, a new feature emerges from an unexpected behavior of the gravitational and inertial mass relation at a very strong gravitational field, with the presence of a divergence that indicates a critical field value. Such a strong gravitational field is just reachable at the early stage of the universe or, perhaps, near massive black holes, and the presence of the critical field value signs that the Equivalence Principle should not be valid at a very strong gravitational field, and certainly it coincides where the quantum effects turns to be relevant^[2].

Hyperbolic motion is the natural generalization of the concept of the Newtonian uniformly accelerated motion due to a constant force applied to a particle, which might be due to an uniform gravitational field. At relativistic level, as the velocity is upper limited by the light velocity, constant force don't imply constant acceleration; instead, it results the above mentioned hyperbolic motion, which denomination comes from the hyperbola that it is drawn in the zt -plane by this kind of motion.

An one dimensional hyperbolic motion of a particle of mass m occurs as a solution of the relativistic equation of motion^{[11], [12]}

$$m \frac{d^2 x^\mu}{d\tau^2} = f^\mu(\tau) , \quad (1)$$

the relativistic force f^μ defined as

$$f^\mu = \gamma \left(\frac{\mathbf{v} \cdot \mathbf{F}}{c}, \mathbf{F} \right) , \quad \gamma = \frac{1}{\sqrt{1 - v^2/c^2}} , \quad (2)$$

when external force $\mathbf{F}$ is parallel to velocity $\mathbf{v}$ and it is constant in the proper referential frame. Supposing the motion along the z -axes, the trajectory is given by

$$(ct, z) = \frac{c^2}{a}(\sinh \lambda\tau, \cosh \lambda\tau), \quad (3)$$

where $a = F/m$ is a constant proper acceleration and $\lambda = a/c$, c the velocity of light. Velocity and acceleration are given by

$$(\dot{z}^0, \dot{z}) = c(\cosh \lambda\tau, \sinh \lambda\tau) \quad (4)$$

and

$$(\ddot{z}^0, \ddot{z}) = a(\sinh \lambda\tau, \cosh \lambda\tau), \quad (5)$$

respectively. The relativistic force responsible by the hyperbolic motion is

$$f^\mu(\tau) = m(\ddot{z}^0, \ddot{z}) = ma(\sinh \lambda\tau, \cosh \lambda\tau). \quad (6)$$

The choice of the metric tensor $g^{\mu\nu}$ is such that $v^\mu v_\mu = -c^2$ for four velocity $v^\mu = \dot{x}^\mu$ and, at non relativistic limit, $a^\mu a_\mu = a^2$ for four acceleration $a^\mu = \ddot{x}^\mu$.

II. HYPERBOLIC MOTION OF CHARGED PARTICLES

Equation of motion of a classical charged particle, including radiation reaction force, is given by the well known Lorentz-Dirac equation

$$ma^\mu(\tau) = f_{ext}^\mu(\tau) + m\tau_0 \left(\dot{a}^\mu - \frac{1}{c^2} a^\nu a_\nu v^\mu \right), \quad (7)$$

where $f_{ext}^\mu(\tau)$ is the external four-force and

$$f_{rad}^\mu(\tau) = m\tau_0 \left(\dot{a}^\mu - \frac{1}{c^2} a^\nu a_\nu v^\mu \right), \quad (8)$$

where

$$\tau_0 = \frac{2}{3} \frac{e^2}{mc^3}, \quad (9)$$

is the Lorentz-Dirac relativistic radiation reaction force. The first term is known as the Schott term ^[5] and it is responsible by the well known non physical runaway solutions. The second is the Rohrlich term, related to the power radiated

$$\mathcal{R} = \frac{dW_{rad}}{dt} = m\tau_0 a^\nu a_\nu. \quad (10)$$

A well known condition for hyperbolic motion is

$$\dot{a}^\mu - \frac{1}{c^2} a^\nu a_\nu v^\mu = 0, \quad (11)$$

which also implies $f_{rad}^\mu(\tau) = 0$, so it seems to be easy to produce hyperbolic motion of charged particles imposing a locally constant external force, as in the uncharged particle case, but it could be a mistake. To have a stable hyperbolic motion we have to get a very sensible balance between external and radiation reaction force, and before it the condition (11) is not true. It means that what happen before is very important to get a stable hyperbolic motion regime and, although we have the same equation (1) after that, the force $f^\mu(\tau)$ is not just $f_{ext}^\mu(\tau)$ anymore. To figure out why, let us consider the Lorentz-Dirac equation written as ^[4]

$$m(1 - \tau_0 \frac{d}{d\tau})a^\mu = f_{ext}^\mu(\tau) - \frac{1}{c^2}\mathcal{R}v^\mu = K^\mu(\tau) . \quad (12)$$

Formal expansion like

$$(1 - \tau_0 \frac{d}{d\tau})^{-1} = 1 + \tau_0 \frac{d}{d\tau} + \tau_0^2 \frac{d^2}{d\tau^2} + \dots \quad (13)$$

enables us to get a formal solution of Lorentz-Dirac equation as

$$ma^\mu(\tau) = \sum_{n=0}^{\infty} \tau_0^n \frac{d^n}{d\tau^n} K^\mu(\tau) . \quad (14)$$

We can insert the mathematical identity

$$\frac{1}{n!} \int_0^\infty s^n e^{-s} ds = 1 \quad (15)$$

to transform (14) in a second order integro-differential equation

$$ma^\mu(\tau) = \int_0^\infty e^{-s} K^\mu(\tau + \tau_0 s) ds , \quad (16)$$

which shows a possible non causal behavior. In an explicit form, we have

$$ma^\mu(\tau) = \int_0^\infty \left(f_{ext}^\mu - \frac{1}{c^2} \mathcal{R}v^\mu \right) \Big|_{\tau + \tau_0 s} e^{-s} ds . \quad (17)$$

Actual hyperbolic motion implies (3-6), from which it is easy to show that $f_{ext}^\mu \rightarrow f^\mu$ and, therefore,

$$\int_0^\infty \left(f^\mu - \frac{1}{c^2} \mathcal{R}v^\mu \right) \Big|_{\tau + \tau_0 s} e^{-s} ds = f^\mu(\tau) , \quad (18)$$

recovering the equation (1), in accordance with equation (7) and condition (11). But, while the motion is approaching the hyperbolic regime, we have a limiting process

$$\dot{a}^\mu - \frac{1}{c^2} a^\nu a_\nu v^\mu \rightarrow 0 \Rightarrow f_{rad}^\mu(\tau) \rightarrow 0 \quad (19)$$

such that the total force behaves to

$$f_{ext}^\mu(\tau) + f_{rad}^\mu(\tau) \rightarrow f^\mu(\tau) . \quad (20)$$

To have an hyperbolic motion, it is imperative that the applied external force goes to

$$f_{ext}^\mu(\tau) \rightarrow f_{ext}^\mu(\tau) = \int_0^\infty f^\mu(\tau + \tau_0 s) e^{-s} ds \quad (21)$$

as well as the radiation reaction force goes to

$$f_{rad}^\mu(\tau) \rightarrow f_{Roh}^\mu(\tau) = \int_0^\infty \frac{1}{c^2} \mathcal{R}v^\mu \Big|_{\tau+\tau_0 s} e^{-s} ds \quad (22)$$

such that, after reaching hyperbolic motion,

$$f^\mu(\tau) = f_{ext}^\mu(\tau) + f_{Roh}^\mu(\tau) , \quad (23)$$

$f^\mu(\tau)$ given by (6).

Using (6), spatial component of equation (21) becomes

$$f_{ext}(\tau) = \frac{ma}{2} \left(\frac{e^{\lambda\tau}}{(1 - \lambda\tau_0)} + \frac{e^{-\lambda\tau}}{(1 + \lambda\tau_0)} \right) \quad (24)$$

and, from equations (5) and (10),

$$f_{Roh}(\tau) = -\frac{ma}{2} \lambda\tau_0 \left(\frac{e^{\lambda\tau}}{(1 - \lambda\tau_0)} - \frac{e^{-\lambda\tau}}{(1 + \lambda\tau_0)} \right) . \quad (25)$$

In the same way, time components become

$$f_{ext}^0(\tau) = \frac{ma}{2} \left(\frac{e^{\lambda\tau}}{(1 - \lambda\tau_0)} - \frac{e^{-\lambda\tau}}{(1 + \lambda\tau_0)} \right) \quad (26)$$

and

$$f_{Roh}^0(\tau) = -\frac{ma}{2} \lambda\tau_0 \left(\frac{e^{\lambda\tau}}{(1 - \lambda\tau_0)} + \frac{e^{-\lambda\tau}}{(1 + \lambda\tau_0)} \right) . \quad (27)$$

From equations (24-27), it is easy to see that the total force (23) satisfies (6) necessary to have an hyperbolic motion.

It shows that the external force necessary to produce an hyperbolic motion of neutral particle,

$$f_{ext}(\tau) = f(\tau) = ma \cosh \lambda\tau , \quad (28)$$

is smaller than the external force (24) necessary to give the same hyperbolic motion of charged particle. All external force applied to neutral particle is used to increase its kinetic energy,

$$\frac{dW}{dt} = vF = mav = mac \sinh \lambda\tau = mc^2 \frac{d\gamma}{dt} . \quad (29)$$

On the other hand, for charged particle, external force (24), that can be written as

$$f_{ext}(\tau) = f(\tau) - f_{Roh}(\tau) , \quad (30)$$

provides the increase of kinetic energy in the same amount as given in (29) and supplies, through $f_{Roh}(\tau)$, the energy lost carried by radiation.

III. UNIFORM GRAVITATIONAL FIELD

We saw in the previous section that charged particle performing hyperbolic motion have to be submitted to an external force given by

$$F_{ext} = \frac{ma}{2} \left(\frac{(1+\beta)}{(1-\lambda\tau_0)} + \frac{(1-\beta)}{(1+\lambda\tau_0)} \right), \quad (31)$$

where $F_{ext}(\tau)$ is the measurable force related to the relativistic force by $f_{ext}(\tau) = \gamma F_{ext}(\tau)$.

The Equivalence Principle [6], [16] says that a particle at rest in the laboratory frame R_{lab} immersed in an uniform gravitational field g is seen by observer in a free falling inertial frame R as performing hyperbolic motion with local constant acceleration $a = g$. The local constant force responsible by its hyperbolic motion is the normal force $F_n = -F_g$ that supports the particle against the gravitational force F_g , so, in absolute value it is equal to mg for uncharged particle. But, for charged particle, the normal force F_n must be equal to the external force F_{ext} of equation (31) for $\beta = 0$,

$$F_{ext} \rightarrow F_n = \frac{mg}{(1-\lambda^2\tau_0^2)}. \quad (32)$$

This result suggests that the observer in the laboratory frame R_{lab} measures the gravitational force acting on charged particle as

$$F_g = \frac{mg}{(1-\lambda^2\tau_0^2)} \quad (33)$$

such that

$$m^* = \frac{m}{(1-\lambda^2\tau_0^2)} \quad (34)$$

should define the gravitational mass m^* of charged particle with inertial mass m . For a typical charged particles as electron or proton, $\tau_0 \simeq 0.62 \times 10^{-23}s$ and $\tau_0 \simeq 0.34 \times 10^{-26}s$, respectively. So, in a field magnitude typical for a terrestrial gravitational field, $g \simeq 10m/s$ and $\lambda = g/c \simeq 3.3 \times 10^{-8}s^{-1}$, we can see that $\lambda^2\tau_0^2 \simeq 4.3 \times 10^{-62}$ and $\lambda^2\tau_0^2 \simeq 2.2 \times 10^{-69}$, respectively for electron and proton, a very small numbers, such that $1 - \lambda^2\tau_0^2 \simeq 1$. As a consequence, the gravitational mass is just slight greater than the inertial mass, $m^* \gtrsim m$, and the gravitational and inertial mass relation defined by equation (34) is much as close to unit, $r_g = m^*/m \simeq 1$. It means that there is no consequence, for practical purpose, due to this slight up deviation of gravitational mass in relation to the inertial mass, at least in a region with gravitational field of magnitude as considered above. In fact, it is true for a very long field interval, starting with $g = 0$ and going until to reach a very strong gravitational field of the order $g \sim 10^{30}m/s^2$, as we are going to show in the next step.

figure

Figure caption: Gravitational and inertial mass relation, r , as function of the gravitational field g . There is a critical point defined by $g_c = c/\tau_0$, which is particle dependent and has the value $g_c \simeq 8.8 \times 10^{36}m/s^2$ for proton.

In a such strong gravitational field region, the field dependence of r_g , see equation (34), starts to manifest, as we can see in the figure 1. It increases very slowly and remains very close to unit, starting with $g = 0$ until to reach the very strong field magnitude of the order $g \sim 10^{30}m/s^2$, approaching the divergence point given by the condition $1 - \lambda^2\tau_0^2 = 0$. Then, r_g increases fast to the infinite as the gravitational field goes to its critical value $g_c = c/\tau_0$. This critical field value is mass dependent, with $g_c \simeq 1.8 \times 10^{35}m/s^2$ and $g_c \simeq 8.8 \times 10^{36}m/s^2$ for electron and proton, respectively. Where is possible to find such strong gravitational field? Certainly at the early time of the Universe and, perhaps, near massive black holes hosted at the center of some galaxies. Using the Newtonian formula of the gravitational field, $g = MG/R^2$, with $M \sim 10^{11}$ times the solar mass, $\sim 1.99 \times 10^{30}kg$. A charged particle must approach as close as $R \approx 1.2 \times 10^{-3}m \approx 1mm$ to see a gravitational field strong as g_c . The matter density to give such strong field is of order $10^{30}kg/m^3$. The divergence of the gravitational and inertial mass relation at the critical field value signs that the Equivalence Principle should not be valid for very strong gravitational field, and the quantum effects turns to be relevant. It lasts some crude questions: what about general relativity at this very strong gravitational field? or is it meaningful the concept of unification of general relativity and quantum theory [?]- [23]?

Above the critical point g_c , the relation r_g turns to be negative, with a possible physical interpretation to be investigated. Perhaps, it should bring us some light to figure out a new physics required to explain the recent data concerned to the cosmic rays with a very high energy, above $10^{19}eV$. There is a possibility that the primary particles of the ultra energetic cosmic rays should be a relic of the charged particles scattered in the early stage of the Universe [24].

IV. CONCLUSION

Although the final equation of motion of neutral and charged particles in hyperbolic motion regime seems to be identical, we realize that there is a fundamental difference between them. For charged particle, actually, the total locally constant force is the sum of an applied and the radiation reaction forces in a subtle combination in such a way that results the same as for neutral particle. An extra force is necessary to supply the same kinetic energy as for uncharged particle plus the energy lost carried by radiation. An interesting implication, as the Equivalence Principle is taken account, is that the gravitational mass of charged particle must be greater than its inertial mass in a very small amount. It is small enough to not be detected by any experimental or practical devices, but it helps us to figure out the condition necessary to get an hyperbolic motion regime and to understand the meaning of its equivalence with a charged particle supported at rest in an uniform gravitational field.

Until now the equality between gravitational and inertial mass was understood as the essence of the Equivalence Principle, condition we realized not to be true for charged particle, where, due to the presence of radiation reaction, a slight deviation of gravitational mass compared to inertial one is necessary to hold the Equivalence Principle. However, a new feature comes from the gravitational and inertial mass relation behavior for a very strong gravitational field, just reachable at the early stage of the universe or, perhaps, near massive black holes. There exists a critical, particle dependent, field value that signs a limit of validity of the Equivalence Principle. It certainly coincides where the quantum effects turns

to be dominant.

REFERENCES

- [1] **P. A. M. Dirac**, *Classical theory of radiating electrons*, Proc. Roy. Soc. **A167**, 148-168 (1938).
- [2] **J. A. Wheeler** and **R. P. Feynman**, *Interaction with the Absorber as the Mechanism of Radiation*, Rev. of Modern Phys. **17**, 157-181 (1945).
- [3] **J. A. Wheeler** and **R. P. Feynman**, *Classical Electrodynamics in Terms of Direct Interparticle Action*, Rev. Mod. Phys. **21**, 425-433 (1949).
- [4] **J. D. Jackson**, *Classical Electrodynamics*, John Wiley & Sons, N.Y. (1962).
- [5] **F. Rohrlich**, *The Equations of Motion of Classical Charges*, Ann. of Physics: **13**, 93-109 (1961).
- [6] **F. Rohrlich**, *Classical Charged Particles*, Addison-Wesley (1965).
- [7] **N. D. Birrell** and **P. C. W. Davies**, *Quantum fields in curved space*, Cambridge Un. Press (1982).
- [8] **Steven Weinberg**, *Gravitation and Cosmology - Principles and Applications of the General Theory of Relativity*, John Willey & Sons, New York (1972).
- [9] **A. A. Logunov** et al, *On incorrect formulations of the equivalence principle*, Physics - Uspekhi Fizicheskikh Nauk **39** (1) 73-79 (1996).
- [10] **V. L. Ginzburg** and **Yu N. Eroshenko**, Comments on the paper by A. A. Logunov et al " *On incorrect formulations of the equivalence principle* " (1996), from electronic site <http://www.ufn.ru/Index96.html>.
- [11] **C. Møller**, *The Theory of Relativity (second edition)*, Oxford University Press (1972).
- [12] **L. Landau** and **E. L. Lifshitz**, *The Classical Theory of Fields*, Pergamon Press, Oxford (1976).
- [13] **C. Teitelboim**, **D. Villarroel** and **Ch. G. Van Weert**, *Classical Electrodynamics of Retarded Fields and Point Particles*, Rev. Nuovo Cimento, v.3,n.9, 1-64 (1980).
- [14] **T. Fulton** and **F. Rohrlich**, *Classical Radiation from a Uniformly Accelerated Charge*, Annals of Phys. **9**, 499-517 (1960).
- [15] **D. G. Boulware**, *Radiation from a Uniformly Accelerated Charge*, Annals of Phys. **124**, 169-188 (1980).
- [16] **F. Rohrlich**, *The Equivalence Principle*, Annals of Phys. **22**, 169-191 (1963).
- [17] See, for example, **C. Itzykson** and **J.-B. Zuber**, *Quantum Field Theory*, McGraw-Hill (1980).
- [18] **M. M. de Souza**, *Classical electrodynamics and the quantum nature of light*, J. Phys. A: Math. Gen. **30**, 6565-6585 (1997).
- [19] **F Denef**, **J. Raeymaekers**, **U. M. Studer** and **W. Troost**, *Classical tunneling as a consequence of radiation reaction forces*, Phys. Rev. **E56**, 3624-3627 (1997).
- [20] **F. Rohrlich**, *The Equivalence Principle Revised*, Found. Phys. **30**: (5) 621-630 (2000).
- [21] **F. Rohrlich**, *Why the principles of inertia and of equivalence hold despite self-interaction*, Phys. Rev. **6312**: (12) 7701- (2001).
- [22] **F. Rohrlich**, *The correct equation of motion of a classical point charge*, Phys. Lett **A283**: (5-6) 276-278 (2001).
- [23] **C. J. Isham**, *Quantum Gravity - An Overview*, *Quantum Gravity 2, a Second Oxford Symposium*, p.1-62 , edited by C. J. Isham, R. Penrose and D. W. Sciama, Clarendon Press, Oxford (1981).

- [24] **M. Nagano, A. A. Watson**, *Observation and implications of the ultrahigh-energy cosmic rays*, Rev. Mod. Phys. 72, 689-732 (2000).

